JOURNAL OF THE AMERICAN PODIATRIC MEDICAL ASSOCIATION

January–December 2011
Volume 101

SUBJECT INDEX

A

Amputation
Acute phase reactants predict the risk of amputation in diabetic foot infection [Akinci et al], 1:1
Human digit partially consumed by a canine during sleep in a patient with neuropathy and diabetes [Rogers and Bevilacqua], 3:275

Anatomy
Anatomical variations of peroneal muscles: a cadaver study in an Indian population and a review of the literature [Prakash et al], 6:505

Anesthesia
Reducing the pain of local 1% lidocaine infiltration with a preceding bacteriostatic saline injection: a double-blind prospective trial [Barrett et al], 3:223

Ankle
Central talar dome lesions: a unique surgical approach with incorporation of a talar allograft for joint reconstitution and restoration of function [Dobbs et al], 2:192
Clinical assessment of ankle joint dorsiflexion: a review of measurement techniques [Gatt and Chockalingam], 1:59
Interrater and intrarater reliability in the measurement of ankle joint dorsiflexion is independent of examiner experience and technique used [Kim et al], 5:407
Management of grade III open dislocated ankle fractures: combined internal fixation with bioabsorbable screws/rods and external fixation [Ye et al], 4:307
Patellofemoral pain syndrome and its association with hip, ankle, and foot function in 16- to 18-year-old high school students: a single-blind case-control study [Mølgaard et al], 3:215
Pigmented villonodular synovitis of the ankle: radiologic characteristics [Hao et al], 3:252
Relationship between the size of bone marrow edema of the talus and ankle pain [Tonbul et al], 5:430
An unusual cause of a cystic lesion with an osteochondral defect in the talus: intraosseous lipoma [Atay et al], 3:269

Antibiotic
In vitro antimicrobial activity of calcium sulfate and hydroxyapatite (Cerament Bone Void Filler) discs using heat-sensitive and non–heat-sensitive antibiotics against methicillin-resistant *Staphylococcus aureus* and *Pseudomonas aeruginosa* [Karr et al], 2:146
Management in the wound-care center outpatient setting of a diabetic patient with forefoot osteomyelitis using Cerament Bone Void Filler impregnated with vancomycin: off-label use [Karr], 3:259
An open-label, three-arm pilot study of the safety and efficacy of topical Microcyn Rx wound care versus oral levofloxacin versus combined therapy for mild diabetic foot infections [Landsman et al], 6:484
Postoperative infection after excisional toenail matrixectomy: a retrospective clinical audit [Rusmir and Salerno], 4:316
What do I do when bone hits the ground? [Brownell], 2:190

Antiretroviral therapy, highly active
Prevalence of plantar verrucae in patients with human immunodeficiency virus infection during the post–highly active antiretroviral therapy era [Johnston et al], 1:35

Apophysitis, calcaneal
Plantar pressures in children with and without Sever's disease [Becerro de Bengoa Vallejo et al], 1:17
Sever's disease: a prospective study of risk factors [Scharfbillig et al], 2:133

Aqueous solution, superoxidized
The use of superoxidized aqueous solution versus saline as a replacement solution in the Versajet lavage system in chronic diabetic foot ulcers: a pilot study [Bowling et al], 2:124

Arch height
Variations in foot posture and mobility between individuals with patellofemoral pain and those in a control group [McPoil et al], 4:289

Arthritis
Dynamic splinting for postoperative hallux limitus: a randomized, controlled trial [John et al], 4:285
Evaluating intratester reliability of manual masking of plantar pressure measurements associated with chronic gout [Rome et al], 5:424
Reliability and validity of the INFOOT three-dimensional foot digitizer for patients with rheumatoid arthritis [De Mits et al], 3:198

Arthroereisis
Stabilization of joint forces of the subtalar complex via HyProCure sinus tarsi stent [Graham et al], 5:390

Arthropathy
The Charcot foot in diabetes [Rogers et al], 5:437

B

Bacteria
In vitro antimicrobial activity of calcium sulfate and hydroxyapatite (Cerament Bone Void Filler) discs using heat-sensitive and non–heat-sensitive antibiotics against methicillin-resistant *Staphylococcus aureus* and *Pseudomonas aeruginosa* [Karr et al], 2:146
Postoperative infection after excisional toenail matrixectomy: a retrospective clinical audit [Rusmir and Salerno], 4:316

Biomechanics
Barefoot running claims and controversies: a review of the literature [Jenkins and Cauthon], 3:231
Changes in the plantar pressure distribution during gait throughout gestation [Ribeiro et al], 5:415
Contralateral peak plantar pressures with a postoperative boot: a preliminary study [Mieras et al], 2:127
Evaluating intratester reliability of manual masking of plantar

JOURNAL OF THE AMERICAN PODIATRIC MEDICAL ASSOCIATION

January–December 2011
Volume 101

SUBJECT INDEX

A

Amputation
Acute phase reactants predict the risk of amputation in diabetic foot infection [Akinci et al], 1:1
Human digit partially consumed by a canine during sleep in a patient with neuropathy and diabetes [Rogers and Bevilacqua], 3:275

Anatomy
Anatomical variations of peroneal muscles: a cadaver study in an Indian population and a review of the literature [Prakash et al], 6:505

Anesthesia
Reducing the pain of local 1% lidocaine infiltration with a preceding bacteriostatic saline injection: a double-blind prospective trial [Barrett et al], 3:223

Ankle
Central talar dome lesions: a unique surgical approach with incorporation of a talar allograft for joint reconstitution and restoration of function [Dobbs et al], 2:192
Clinical assessment of ankle joint dorsiflexion: a review of measurement techniques [Gatt and Chockalingam], 1:59
Interrater and intrarater reliability in the measurement of ankle joint dorsiflexion is independent of examiner experience and technique used [Kim et al], 5:407
Management of grade III open dislocated ankle fractures: combined internal fixation with bioabsorbable screws/rods and external fixation [Ye et al], 4:307
Patellofemoral pain syndrome and its association with hip, ankle, and foot function in 16- to 18-year-old high school students: a single-blind case-control study [Mølgaard et al], 3:215
Pigmented villonodular synovitis of the ankle: radiologic characteristics [Hao et al], 3:252
Relationship between the size of bone marrow edema of the talus and ankle pain [Tonbul et al], 5:430
An unusual cause of a cystic lesion with an osteochondral defect in the talus: intraosseous lipoma [Atay et al], 3:269

Antibiotic
In vitro antimicrobial activity of calcium sulfate and hydroxyapatite (Cerament Bone Void Filler) discs using heat-sensitive and non–heat-sensitive antibiotics against methicillin-resistant *Staphylococcus aureus* and *Pseudomonas aeruginosa* [Karr et al], 2:146
Management in the wound-care center outpatient setting of a diabetic patient with forefoot osteomyelitis using Cerament Bone Void Filler impregnated with vancomycin: off-label use [Karr], 3:259
An open-label, three-arm pilot study of the safety and efficacy of topical Microcyn Rx wound care versus oral levofloxacin versus combined therapy for mild diabetic foot infections [Landsman et al], 6:484
Postoperative infection after excisional toenail matrixectomy: a retrospective clinical audit [Rusmir and Salerno], 4:316
What do I do when bone hits the ground? [Brownell], 2:190

Antiretroviral therapy, highly active
Prevalence of plantar verrucae in patients with human immunodeficiency virus infection during the post–highly active antiretroviral therapy era [Johnston et al], 1:35

Apophysitis, calcaneal
Plantar pressures in children with and without Sever's disease [Becerro de Bengoa Vallejo et al], 1:17
Sever's disease: a prospective study of risk factors [Scharfbillig et al], 2:133

Aqueous solution, superoxidized
The use of superoxidized aqueous solution versus saline as a replacement solution in the Versajet lavage system in chronic diabetic foot ulcers: a pilot study [Bowling et al], 2:124

Arch height
Variations in foot posture and mobility between individuals with patellofemoral pain and those in a control group [McPoil et al], 4:289

Arthritis
Dynamic splinting for postoperative hallux limitus: a randomized, controlled trial [John et al], 4:285
Evaluating intratester reliability of manual masking of plantar pressure measurements associated with chronic gout [Rome et al], 5:424
Reliability and validity of the INFOOT three-dimensional foot digitizer for patients with rheumatoid arthritis [De Mits et al], 3:198

Arthroereisis
Stabilization of joint forces of the subtalar complex via HyProCure sinus tarsi stent [Graham et al], 5:390

Arthropathy
The Charcot foot in diabetes [Rogers et al], 5:437

B

Bacteria
In vitro antimicrobial activity of calcium sulfate and hydroxyapatite (Cerament Bone Void Filler) discs using heat-sensitive and non–heat-sensitive antibiotics against methicillin-resistant *Staphylococcus aureus* and *Pseudomonas aeruginosa* [Karr et al], 2:146
Postoperative infection after excisional toenail matrixectomy: a retrospective clinical audit [Rusmir and Salerno], 4:316

Biomechanics
Barefoot running claims and controversies: a review of the literature [Jenkins and Cauthon], 3:231
Changes in the plantar pressure distribution during gait throughout gestation [Ribeiro et al], 5:415
Contralateral peak plantar pressures with a postoperative boot: a preliminary study [Mieras et al], 2:127
Evaluating intratester reliability of manual masking of plantar

pressure measurements associated with chronic gout [Rome et al], 5:424
Foot biomechanics in patients with diabetes mellitus: doubts regarding the relationship between neuropathy, foot motion, and deformities [Lázaro-Martínez et al], 3:208
Foot orthoses: a review focusing on kinematics [Chevalier and Chockalingam], 4:341
Hallux limitus and its relationship with the internal rotational pattern of the lower limb [Munuera et al], 6:467
In-shoe plantar pressure distribution in nonneuropathic type 2 diabetic patients in Singapore [Tong et al], 6:509
The in vitro reliability of the CODA MPX30 as the basis for a method of assessing the in vivo motion of the subtalar joint [Birch and Deschamps], 5:400
The medial oblique shell inclination technique: a method to increase subtalar supination moments in foot orthoses [Harradine et al], 6:523
Plantar pressures in children with and without Sever's disease [Becerro de Bengoa Vallejo et al], 1:17
Quantification of skin marker movement at the malleoli and talar heads [Birch and Deschamps], 6:497
Sever's disease: a prospective study of risk factors [Scharfbillig et al], 2:133
Soft-tissue movement at the foot during the stance phase of walking [Chen et al], 1:25
Stabilization of joint forces of the subtalar complex via HyProCure sinus tarsi stent [Graham et al], 5:390

Bone
Bilateral tibial hallux sesamoid agenesis and fibular hallux sesamoid hypoplasia in a patient with bilateral hallux valgus [Marinozzi et al], 5:452
Central talar dome lesions: a unique surgical approach with incorporation of a talar allograft for joint reconstitution and restoration of function [Dobbs et al], 2:192
The Charcot foot in diabetes [Rogers et al], 5:437
Foot salvage after loss of the first and second metatarsal rays with a free fibular osteocutaneous flap [Salgado et al], 6:531
Freiberg's disease: quantitative assessment of osteonecrosis on three-dimensional CT [Chun et al], 4:335
In vitro antimicrobial activity of calcium sulfate and hydroxyapatite (Cerament Bone Void Filler) discs using heat-sensitive and non–heat-sensitive antibiotics against methicillin-resistant *Staphylococcus aureus* and *Pseudomonas aeruginosa* [Karr et al], 2:146
Management in the wound-care center outpatient setting of a diabetic patient with forefoot osteomyelitis using Cerament Bone Void Filler impregnated with vancomycin: off-label use [Karr], 3:259
Metastatic renal cell carcinoma presenting as foot metastasis: case report and review of the literature [Choufani et al], 3:265
Navicular stress reactions in runners: a review of evaluation and management of a competitive athlete [Yoho and Wells], 5:447
Relationship between the size of bone marrow edema of the talus and ankle pain [Tonbul et al], 5:430
Surgical treatment of diaphyseal stress fractures of the fifth metatarsal in competitive athletes: long-term follow-up and computerized pedobarographic analysis [Smoljanovic et al], 6:517
The symptomatic os vesalianum as an uncommon cause of lateral foot pain: a case report [Wilson et al], 4:356
An unusual cause of a cystic lesion with an osteochondral defect in the talus: intraosseous lipoma [Atay et al], 3:269
What do I do when bone hits the ground? [Brownell], 2:190

Braces, nail
A quantitative method for measuring forces applied by nail braces [Erdogan], 3:247

Bunionectomy
What do I do when bone hits the ground? [Brownell], 2:190

C

Calcaneus
Plantar pressures in children with and without Sever's disease [Becerro de Bengoa Vallejo et al], 1:17
Sever's disease: a prospective study of risk factors [Scharfbillig et al], 2:133
Surgical treatment of pressure ulcers of the heel in skilled nursing facilities: a 12-year retrospective study of 57 patients [Han and Ezquerro], 2:167

Calcium sulfate
In vitro antimicrobial activity of calcium sulfate and hydroxyapatite (Cerament Bone Void Filler) discs using heat-sensitive and non–heat-sensitive antibiotics against methicillin-resistant *Staphylococcus aureus* and *Pseudomonas aeruginosa* [Karr et al], 2:146
Management in the wound-care center outpatient setting of a diabetic patient with forefoot osteomyelitis using Cerament Bone Void Filler impregnated with vancomycin: off-label use [Karr], 3:259

Callus
Foot biomechanics in patients with diabetes mellitus: doubts regarding the relationship between neuropathy, foot motion, and deformities [Lázaro-Martínez et al], 3:208

Cancer
Invasive cutaneous squamous cell carcinoma presenting as a nonhealing radiation wound: a case report [Longobardi et al], 4:360
Metastatic renal cell carcinoma presenting as foot metastasis: case report and review of the literature [Choufani et al], 3:265
Small blue round cell tumor: an unusual case presentation in the foot [Devall et al], 4:363

Canine
Human digit partially consumed by a canine during sleep in a patient with neuropathy and diabetes [Rogers and Bevilacqua], 3:275

Carcinoma
Invasive cutaneous squamous cell carcinoma presenting as a nonhealing radiation wound: a case report [Longobardi et al], 4:360
Metastatic renal cell carcinoma presenting as foot metastasis: case report and review of the literature [Choufani et al], 3:265

Cerament Bone Void Filler
In vitro antimicrobial activity of calcium sulfate and hydroxyapatite (Cerament Bone Void Filler) discs using heat-sensitive and non–heat-sensitive antibiotics against methicillin-resistant *Staphylococcus aureus* and *Pseudomonas aeruginosa* [Karr et al], 2:146
Management in the wound-care center outpatient setting of a diabetic patient with forefoot osteomyelitis using Cerament Bone Void Filler impregnated with vancomycin: off-label use [Karr], 3:259

Charcot-Marie-Tooth disease
Charcot-Marie-Tooth disease complicating type 2 diabetes [Win et al], 4:349

Charcot's foot
The Charcot foot in diabetes [Rogers et al], 5:437

Chigoe
Tungiasis in a beach volleyball player: a case report [Veraldi et al], 4:353

Chondrocalcinosis
Mimicry in older patients: tophaceous pseudogout as a tumorlike lesion: a case report [Erhardt et al], 5:462

Claw toe
Dorsal approach to transfer of the flexor digitorum brevis tendon [Becerro de Bengoa Vallejo et al], 4:297

CODA MPX30 system
The in vitro reliability of the CODA MPX30 as the basis for a method of assessing the in vivo motion of the subtalar joint [Birch and Deschamps], 5:400

Computer technology
The use of focused electronic medical record forms to improve health-care outcomes [Caldwell et al], 4:331

Congenital deformity
Lengthening of the midfoot in a case of lower-extremity hemimelia [Kraus et al], 5:456

Contracture
Surgical treatment of pressure ulcers of the heel in skilled nursing facilities: a 12-year retrospective study of 57 patients [Han and Ezquerro], 2:167

Costs, health-care
The economic value of specialized lower-extremity medical care by podiatric physicians in the treatment of diabetic foot ulcers [Carls et al], 2:93

D

Debridement
The use of superoxidized aqueous solution versus saline as a replacement solution in the Versajet lavage system in chronic diabetic foot ulcers: a pilot study [Bowling et al], 2:124

Diabetes mellitus
Acute phase reactants predict the risk of amputation in diabetic foot infection [Akinci et al], 1:1
Autologous skin fibroblast and keratinocyte grafts in the treatment of chronic foot ulcers in aging type 2 diabetic patients [Monami et al], 1:55
The Charcot foot in diabetes [Rogers et al], 5:437
Charcot-Marie-Tooth disease complicating type 2 diabetes [Win et al], 4:349
A diabetic foot ulcer pilot study: does silicone gel sheeting reduce the incidence of reulceration? [Westphal et al], 2:116
The economic value of specialized lower-extremity medical care by podiatric physicians in the treatment of diabetic foot ulcers [Carls et al], 2:93
Foot biomechanics in patients with diabetes mellitus: doubts regarding the relationship between neuropathy, foot motion, and deformities [Lázaro-Martínez et al], 3:208
Foot care recommendations: an excerpt from *Diabetes Care* [American Diabetes Association], 1:75
Human digit partially consumed by a canine during sleep in a patient with neuropathy and diabetes [Rogers and Bevilacqua], 3:275
In-shoe plantar pressure distribution in nonneuropathic type 2 diabetic patients in Singapore [Tong et al], 6:509
Management in the wound-care center outpatient setting of a diabetic patient with forefoot osteomyelitis using Cerament Bone Void Filler impregnated with vancomycin: off-label use [Karr], 3:259
Motivational interviewing by podiatric physicians: a method for improving patient self-care of the diabetic foot [Gabbay et al], 1:78
An open-label, three-arm pilot study of the safety and efficacy of topical Microcyn Rx wound care versus oral levofloxacin versus combined therapy for mild diabetic foot infections [Landsman et al], 6:484
Prevalence of toenail onychomycosis in patients with type 2 diabetes mellitus and evaluation of risk factors [Gulcan et al], 1:49
The use of superoxidized aqueous solution versus saline as a replacement solution in the Versajet lavage system in chronic diabetic foot ulcers: a pilot study [Bowling et al], 2:124

Dorsiflexion
Clinical assessment of ankle joint dorsiflexion: a review of measurement techniques [Gatt and Chockalingam], 1:59
Dynamic splinting for postoperative hallux limitus: a randomized, controlled trial [John et al], 4:285
Hallux limitus and its relationship with the internal rotational pattern of the lower limb [Munuera et al], 6:467
Interrater and intrarater reliability in the measurement of ankle joint dorsiflexion is independent of examiner experience and technique used [Kim et al], 5:407

E

Edema
Relationship between the size of bone marrow edema of the talus and ankle pain [Tonbul et al], 5:430

Education
Effect of an educational intervention on nurses' knowledge of foot care and on the foot health of older residents [Stolt et al], 2:159

Electronic medical records
The use of focused electronic medical record forms to improve health-care outcomes [Caldwell et al], 4:331

Equinus deformity
Clinical assessment of ankle joint dorsiflexion: a review of measurement techniques [Gatt and Chockalingam], 1:59
Interrater and intrarater reliability in the measurement of ankle joint dorsiflexion is independent of examiner experience and technique used [Kim et al], 5:407
Plantar pressures in children with and without Sever's disease [Becerro de Bengoa Vallejo et al], 1:17

Evidence-based medicine
The use of focused electronic medical record forms to improve health-care outcomes [Caldwell et al], 4:331

F

Fasciitis, plantar
The correlation between plantar fascia thickness and symptoms of plantar fasciitis [Mahowald et al], 5:385

Femur
Effect of foot orthoses contour on pain perception in individuals with patellofemoral pain [McPoil et al], 1:7
Patellofemoral pain syndrome and its association with hip, ankle, and foot function in 16- to 18-year-old high school students: a single-blind case-control study [Mølgaard et al], 3:215
Variations in foot posture and mobility between individuals with patellofemoral pain and those in a control group [McPoil et al], 4:289

Fibromatosis
Extra-abdominal desmoid fibromatosis in the foot: a case study [MacGill et al], 1:70

Fibula
Foot salvage after loss of the first and second metatarsal rays with a free fibular osteocutaneous flap [Salgado et al], 6:531

Flap
Distally based adipofascial sural flap for foot and ankle reconstruction [Mojallal et al], 1:41

Foot salvage after loss of the first and second metatarsal rays with a free fibular osteocutaneous flap [Salgado et al], 6:531

Flatfoot
Posterior tibial tendon dysfunction: a review [Durrant et al], 2:176

Flea
Tungiasis in a beach volleyball player: a case report [Veraldi et al], 4:353

Foot care
Effect of an educational intervention on nurses' knowledge of foot care and on the foot health of older residents [Stolt et al], 2:159
Foot care recommendations: an excerpt from *Diabetes Care*, 1:75
Motivational interviewing by podiatric physicians: a method for improving patient self-care of the diabetic foot [Gabbay et al], 1:78

Foot posture
Patellofemoral pain syndrome and its association with hip, ankle, and foot function in 16- to 18-year-old high school students: a single-blind case-control study [Mølgaard et al], 3:215
Reliability and validity of the INFOOT three-dimensional foot digitizer for patients with rheumatoid arthritis [De Mits et al], 3:198
Variations in foot posture and mobility between individuals with patellofemoral pain and those in a control group [McPoil et al], 4:289

Foot structure
Reliability and validity of the INFOOT three-dimensional foot digitizer for patients with rheumatoid arthritis [De Mits et al], 3:198

Footwear
Barefoot running claims and controversies: a review of the literature [Jenkins and Cauthon], 3:231
Contralateral peak plantar pressures with a postoperative boot: a preliminary study [Mieras et al], 2:127

Forces
A quantitative method for measuring forces applied by nail braces [Erdogan], 3:247
Stabilization of joint forces of the subtalar complex via HyProCure sinus tarsi stent [Graham et al], 5:390

Forefoot
Management in the wound-care center outpatient setting of a diabetic patient with forefoot osteomyelitis using Cerament Bone Void Filler impregnated with vancomycin: off-label use [Karr], 3:259

Fracture
Foot salvage after loss of the first and second metatarsal rays with a free fibular osteocutaneous flap [Salgado et al], 6:531
Navicular stress reactions in runners: a review of evaluation and management of a competitive athlete [Yoho and Wells], 5:447
Surgical treatment of diaphyseal stress fractures of the fifth metatarsal in competitive athletes: long-term follow-up and computerized pedobarographic analysis [Smoljanovic et al], 6:517

Freiberg's disease
Freiberg's disease: quantitative assessment of osteonecrosis on three-dimensional CT [Chun et al], 4:335

G

Gait
Barefoot running claims and controversies: a review of the literature [Jenkins and Cauthon], 3:231
Changes in the plantar pressure distribution during gait throughout gestation [Ribeiro et al], 5:415
Evaluating intratester reliability of manual masking of plantar pressure measurements associated with chronic gout [Rome et al], 5:424
Foot orthoses: a review focusing on kinematics [Chevalier and Chockalingam], 4:341
The in vitro reliability of the CODA MPX30 as the basis for a method of assessing the in vivo motion of the subtalar joint [Birch and Deschamps], 5:400
Quantification of skin marker movement at the malleoli and talar heads [Birch and Deschamps], 6:497
Soft-tissue movement at the foot during the stance phase of walking [Chen et al], 1:25
Stabilization of joint forces of the subtalar complex via HyProCure sinus tarsi stent [Graham et al], 5:390

Gastrocnemius soleus complex
Plantar pressures in children with and without Sever's disease [Becerro de Bengoa Vallejo et al], 1:17

Geriatric population
Autologous skin fibroblast and keratinocyte grafts in the treatment of chronic foot ulcers in aging type 2 diabetic patients [Monami et al], 1:55
Effect of an educational intervention on nurses' knowledge of foot care and on the foot health of older residents [Stolt et al], 2:159
Mimicry in older patients: tophaceous pseudogout as a tumorlike lesion: a case report [Erhardt et al], 5:462

Gestation
Changes in the plantar pressure distribution during gait throughout gestation [Ribeiro et al], 5:415

Goniometry
Interrater and intrarater reliability in the measurement of ankle joint dorsiflexion is independent of examiner experience and technique used [Kim et al], 5:407

Gout
Evaluating intratester reliability of manual masking of plantar pressure measurements associated with chronic gout [Rome et al], 5:424

Graft
Autologous skin fibroblast and keratinocyte grafts in the treatment of chronic foot ulcers in aging type 2 diabetic patients [Monami et al], 1:55
Central talar dome lesions: a unique surgical approach with incorporation of a talar allograft for joint reconstitution and restoration of function [Dobbs et al], 2:192
Foot salvage after loss of the first and second metatarsal rays with a free fibular osteocutaneous flap [Salgado et al], 6:531

Guidelines, authors'
Guidelines for authors [Joseph], 1:85, 4:371

H

Hallux
Bilateral tibial hallux sesamoid agenesis and fibular hallux sesamoid hypoplasia in a patient with bilateral hallux valgus [Marinozzi et al], 5:452
Dynamic splinting for postoperative hallux limitus: a randomized, controlled trial [John et al], 4:285
Hallux limitus and its relationship with the internal rotational pattern of the lower limb [Munuera et al], 6:467
Human digit partially consumed by a canine during sleep in a patient with neuropathy and diabetes [Rogers and Bevilacqua], 3:275

Hallux limitus
Dynamic splinting for postoperative hallux limitus: a randomized, controlled trial [John et al], 4:285
Hallux limitus and its relationship with the internal rotational pattern of the lower limb [Munuera et al], 6:467

Hallux valgus
Bilateral tibial hallux sesamoid agenesis and fibular hallux sesamoid hypoplasia in a patient with bilateral hallux valgus [Marinozzi et al], 5:452

Hammer toe
Dorsal approach to transfer of the flexor digitorum brevis tendon [Becerro de Bengoa Vallejo et al], 4:297

Heel
The correlation between plantar fascia thickness and symptoms of plantar fasciitis [Mahowald et al], 5:385
Surgical treatment of pressure ulcers of the heel in skilled nursing facilities: a 12-year retrospective study of 57 patients [Han and Ezquerro], 2:167

Hemimelia
Lengthening of the midfoot in a case of lower-extremity hemimelia [Kraus et al], 5:456

Hip
Patellofemoral pain syndrome and its association with hip, ankle, and foot function in 16- to 18-year-old high school students: a single-blind case-control study [Mølgaard et al], 3:215

Human immunodeficiency virus
Prevalence of plantar verrucae in patients with human immunodeficiency virus infection during the post–highly active antiretroviral therapy era [Johnston et al], 1:35

Hydroxyapatite
In vitro antimicrobial activity of calcium sulfate and hydroxyapatite (Cerament Bone Void Filler) discs using heat-sensitive and non–heat-sensitive antibiotics against methicillin-resistant *Staphylococcus aureus* and *Pseudomonas aeruginosa* [Karr et al], 2:146
Management in the wound-care center outpatient setting of a diabetic patient with forefoot osteomyelitis using Cerament Bone Void Filler impregnated with vancomycin: off-label use [Karr], 3:259

HyProCure stent
Stabilization of joint forces of the subtalar complex via HyProCure sinus tarsi stent [Graham et al], 5:390

I

Ilizarov fixator
Lengthening of the midfoot in a case of lower-extremity hemimelia [Kraus et al], 5:456

Imaging
The correlation between plantar fascia thickness and symptoms of plantar fasciitis [Mahowald et al], 5:385
Freiberg's disease: quantitative assessment of osteonecrosis on three-dimensional CT [Chun et al], 4:335
Pigmented villonodular synovitis of the ankle: radiologic characteristics [Hao et al], 3:252
Reliability and validity of the INFOOT three-dimensional foot digitizer for patients with rheumatoid arthritis [De Mits et al], 3:198
Validation of the talar–second metatarsal angle as a standard measurement for radiographic evaluation [Graham et al], 6:475

Infection
Acute phase reactants predict the risk of amputation in diabetic foot infection [Akinci et al], 1:1
In vitro antimicrobial activity of calcium sulfate and hydroxyapatite (Cerament Bone Void Filler) discs using heat-sensitive and non–heat-sensitive antibiotics against methicillin-resistant *Staphylococcus aureus* and *Pseudomonas aeruginosa* [Karr et al], 2:146
Management in the wound-care center outpatient setting of a diabetic patient with forefoot osteomyelitis using Cerament Bone Void Filler impregnated with vancomycin: off-label use [Karr], 3:259
An open-label, three-arm pilot study of the safety and efficacy of topical Microcyn Rx wound care versus oral levofloxacin versus combined therapy for mild diabetic foot infections [Landsman et al], 6:484
Postoperative infection after excisional toenail matrixectomy: a retrospective clinical audit [Rusmir and Salerno], 4:316
Prevalence of plantar verrucae in patients with human immunodeficiency virus infection during the post–highly active antiretroviral therapy era [Johnston et al], 1:35
Prevalence of toenail onychomycosis in patients with type 2 diabetes mellitus and evaluation of risk factors [Gulcan et al], 1:49

Injection
Reducing the pain of local 1% lidocaine infiltration with a preceding bacteriostatic saline injection: a double-blind prospective trial [Barrett et al], 3:223

Injury
Barefoot running claims and controversies: a review of the literature [Jenkins and Cauthon], 3:231
Development of a novel rating system to assess lower-limb comfort [Kinchington et al], 5:371
Foot salvage after loss of the first and second metatarsal rays with a free fibular osteocutaneous flap [Salgado et al], 6:531
Human digit partially consumed by a canine during sleep in a patient with neuropathy and diabetes [Rogers and Bevilacqua], 3:275
Management of grade III open dislocated ankle fractures: combined internal fixation with bioabsorbable screws/rods and external fixation [Ye et al], 4:307
Navicular stress reactions in runners: a review of evaluation and management of a competitive athlete [Yoho and Wells], 5:447
Surgical treatment of diaphyseal stress fractures of the fifth metatarsal in competitive athletes: long-term follow-up and computerized pedobarographic analysis [Smoljanovic et al], 6:517

Insect
Tungiasis in a beach volleyball player: a case report [Veraldi et al], 4:353

Interviewing, motivational
Motivational interviewing by podiatric physicians: a method for improving patient self-care of the diabetic foot [Gabbay et al], 1:78

Ischemia
Relationship between the size of bone marrow edema of the talus and ankle pain [Tonbul et al], 5:430

J

Joint
Bilateral tibial hallux sesamoid agenesis and fibular hallux sesamoid hypoplasia in a patient with bilateral hallux valgus [Marinozzi et al], 5:452
Central talar dome lesions: a unique surgical approach with incorporation of a talar allograft for joint reconstitution and restoration of function [Dobbs et al], 2:192

The Charcot foot in diabetes [Rogers et al], 5:437
Clinical assessment of ankle joint dorsiflexion: a review of measurement techniques [Gatt and Chockalingam], 1:59
Dynamic splinting for postoperative hallux limitus: a randomized, controlled trial [John et al], 4:285
Effect of foot orthoses contour on pain perception in individuals with patellofemoral pain [McPoil et al], 1:7
Evaluating intratester reliability of manual masking of plantar pressure measurements associated with chronic gout [Rome et al], 5:424
Foot biomechanics in patients with diabetes mellitus: doubts regarding the relationship between neuropathy, foot motion, and deformities [Lázaro-Martínez et al], 3:208
Hallux limitus and its relationship with the internal rotational pattern of the lower limb [Munuera et al], 6:467
Interrater and intrarater reliability in the measurement of ankle joint dorsiflexion is independent of examiner experience and technique used [Kim et al], 5:407
The in vitro reliability of the CODA MPX30 as the basis for a method of assessing the in vivo motion of the subtalar joint [Birch and Deschamps], 5:400
Management of grade III open dislocated ankle fractures: combined internal fixation with bioabsorbable screws/rods and external fixation [Ye et al], 4:307
The medial oblique shell inclination technique: a method to increase subtalar supination moments in foot orthoses [Harradine et al], 6:523
Patellofemoral pain syndrome and its association with hip, ankle, and foot function in 16- to 18-year-old high school students: a single-blind case-control study [Mølgaard et al], 3:215
Pigmented villonodular synovitis of the ankle: radiologic characteristics [Hao et al], 3:252
Quantification of skin marker movement at the malleoli and talar heads [Birch and Deschamps], 6:497
Reliability and validity of the INFOOT three-dimensional foot digitizer for patients with rheumatoid arthritis [De Mits et al], 3:198
Stabilization of joint forces of the subtalar complex via HyProCure sinus tarsi stent [Graham et al], 5:390

K

Kidney
Metastatic renal cell carcinoma presenting as foot metastasis: case report and review of the literature [Choufani et al], 3:265

Kinematics
Foot orthoses: a review focusing on kinematics [Chevalier and Chockalingam], 4:341
Quantification of skin marker movement at the malleoli and talar heads [Birch and Deschamps], 6:497
Soft-tissue movement at the foot during the stance phase of walking [Chen et al], 1:25
Stabilization of joint forces of the subtalar complex via HyProCure sinus tarsi stent [Graham et al], 5:390

Knee
Effect of foot orthoses contour on pain perception in individuals with patellofemoral pain [McPoil et al], 1:7
Patellofemoral pain syndrome and its association with hip, ankle, and foot function in 16- to 18-year-old high school students: a single-blind case-control study [Mølgaard et al], 3:215
Variations in foot posture and mobility between individuals with patellofemoral pain and those in a control group [McPoil et al], 4:289

L

Lactic acid
Clinical evaluation of 35% urea in a water-lipid–based foam containing lactic acid for treatment of mild-to-moderate xerosis of the foot [Grossman], 2:153

Lavage
The use of superoxidized aqueous solution versus saline as a replacement solution in the Versajet lavage system in chronic diabetic foot ulcers: a pilot study [Bowling et al], 2:124

Lesion
Central talar dome lesions: a unique surgical approach with incorporation of a talar allograft for joint reconstitution and restoration of function [Dobbs et al], 2:192
Extra-abdominal desmoid fibromatosis in the foot: a case study [MacGill et al], 1:70
Giant cell tumor of the flexor hallucis longus tendon sheath: a case study [Findling et al], 2:187
Invasive cutaneous squamous cell carcinoma presenting as a nonhealing radiation wound: a case report [Longobardi et al], 4:360
Metastatic renal cell carcinoma presenting as foot metastasis: case report and review of the literature [Choufani et al], 3:265
Mimicry in older patients: tophaceous pseudogout as a tumorlike lesion: a case report [Erhardt et al], 5:462
Pigmented villonodular synovitis of the ankle: radiologic characteristics [Hao et al], 3:252
Small blue round cell tumor: an unusual case presentation in the foot [Devall et al], 4:363
An unusual cause of a cystic lesion with an osteochondral defect in the talus: intraosseous lipoma [Atay et al], 3:269

Levofloxacin
An open-label, three-arm pilot study of the safety and efficacy of topical Microcyn Rx wound care versus oral levofloxacin versus combined therapy for mild diabetic foot infections [Landsman et al], 6:484

Lidocaine
Reducing the pain of local 1% lidocaine infiltration with a preceding bacteriostatic saline injection: a double-blind prospective trial [Barrett et al], 3:223

Lipoma
An unusual cause of a cystic lesion with an osteochondral defect in the talus: intraosseous lipoma [Atay et al], 3:269

M

Malleolus
Quantification of skin marker movement at the malleoli and talar heads [Birch and Deschamps], 6:497

Matrixectomy
Postoperative infection after excisional toenail matrixectomy: a retrospective clinical audit [Rusmir and Salerno], 4:316

Measurement
Clinical assessment of ankle joint dorsiflexion: a review of measurement techniques [Gatt and Chockalingam], 1:59
Development of a novel rating system to assess lower-limb comfort [Kinchington et al], 5:371
Evaluating intratester reliability of manual masking of plantar pressure measurements associated with chronic gout [Rome et al], 5:424
Freiberg's disease: quantitative assessment of osteonecrosis on three-dimensional CT [Chun et al], 4:335
Interrater and intrarater reliability in the measurement of ankle joint dorsiflexion is independent of examiner experience and technique used [Kim et al], 5:407
The in vitro reliability of the CODA MPX30 as the basis for a method of assessing the in vivo motion of the subtalar joint [Birch and Deschamps], 5:400
Quantification of skin marker movement at the malleoli and talar heads [Birch and Deschamps], 6:497
A quantitative method for measuring forces applied by nail braces [Erdogan], 3:247

Reliability and validity of the INFOOT three-dimensional foot digitizer for patients with rheumatoid arthritis [De Mits et al], 3:198
Validation of the talar–second metatarsal angle as a standard measurement for radiographic evaluation [Graham et al], 6:475

Metatarsal
Dynamic splinting for postoperative hallux limitus: a randomized, controlled trial [John et al], 4:285
Foot salvage after loss of the first and second metatarsal rays with a free fibular osteocutaneous flap [Salgado et al], 6:531
Freiberg's disease: quantitative assessment of osteonecrosis on three-dimensional CT [Chun et al], 4:335
Surgical treatment of diaphyseal stress fractures of the fifth metatarsal in competitive athletes: long-term follow-up and computerized pedobarographic analysis [Smoljanovic et al], 6:517
Validation of the talar–second metatarsal angle as a standard measurement for radiographic evaluation [Graham et al], 6:475
What do I do when bone hits the ground? [Brownell], 2:190

Microcyn Rx
An open-label, three-arm pilot study of the safety and efficacy of topical Microcyn Rx wound care versus oral levofloxacin versus combined therapy for mild diabetic foot infections [Landsman et al], 6:484

Midfoot
Lengthening of the midfoot in a case of lower-extremity hemimelia [Kraus et al], 5:456

Mimicry
Mimicry in older patients: tophaceous pseudogout as a tumorlike lesion: a case report [Erhardt et al], 5:462

Muscle
Anatomical variations of peroneal muscles: a cadaver study in an Indian population and a review of the literature [Prakash et al], 6:505

N

Nail
Postoperative infection after excisional toenail matrixectomy: a retrospective clinical audit [Rusmir and Salerno], 4:316
Prevalence of toenail onychomycosis in patients with type 2 diabetes mellitus and evaluation of risk factors [Gulcan et al], 1:49
A quantitative method for measuring forces applied by nail braces [Erdogan], 3:247

Navicular
Navicular stress reactions in runners: a review of evaluation and management of a competitive athlete [Yoho and Wells], 5:447

Necrosis
Freiberg's disease: quantitative assessment of osteonecrosis on three-dimensional CT [Chun et al], 4:335

Neuropathy
The Charcot foot in diabetes [Rogers et al], 5:437
Charcot-Marie-Tooth disease complicating type 2 diabetes [Win et al], 4:349
Foot biomechanics in patients with diabetes mellitus: doubts regarding the relationship between neuropathy, foot motion, and deformities [Lázaro-Martínez et al], 3:208
Human digit partially consumed by a canine during sleep in a patient with neuropathy and diabetes [Rogers and Bevilacqua], 3:275
In-shoe plantar pressure distribution in nonneuropathic type 2 diabetic patients in Singapore [Tong et al], 6:509

Nursing
Effect of an educational intervention on nurses' knowledge of foot care and on the foot health of older residents [Stolt et al], 2:159
Surgical treatment of pressure ulcers of the heel in skilled nursing facilities: a 12-year retrospective study of 57 patients [Han and Ezquerro], 2:167

O

Onychocryptosis
Postoperative infection after excisional toenail matrixectomy: a retrospective clinical audit [Rusmir and Salerno], 4:316
A quantitative method for measuring forces applied by nail braces [Erdogan], 3:247

Onychomycosis
Prevalence of toenail onychomycosis in patients with type 2 diabetes mellitus and evaluation of risk factors [Gulcan et al], 1:49

Orthoses
Effect of foot orthoses contour on pain perception in individuals with patellofemoral pain [McPoil et al], 1:7
Foot orthoses: a review focusing on kinematics [Chevalier and Chockalingam], 4:341
The medial oblique shell inclination technique: a method to increase subtalar supination moments in foot orthoses [Harradine et al], 6:523

Osteoarthropathy
The Charcot foot in diabetes [Rogers et al], 5:437

Osteochondritis dissecans
Central talar dome lesions: a unique surgical approach with incorporation of a talar allograft for joint reconstitution and restoration of function [Dobbs et al], 2:192

Osteomyelitis
In vitro antimicrobial activity of calcium sulfate and hydroxyapatite (Cerament Bone Void Filler) discs using heat-sensitive and non–heat-sensitive antibiotics against methicillin-resistant *Staphylococcus aureus* and *Pseudomonas aeruginosa* [Karr et al], 2:146
Management in the wound-care center outpatient setting of a diabetic patient with forefoot osteomyelitis using Cerament Bone Void Filler impregnated with vancomycin: off-label use [Karr], 3:259

Osteonecrosis
Freiberg's disease: quantitative assessment of osteonecrosis on three-dimensional CT [Chun et al], 4:335

Os vesalianum pedis
The symptomatic os vesalianum as an uncommon cause of lateral foot pain: a case report [Wilson et al], 4:356

P

Pain
The correlation between plantar fascia thickness and symptoms of plantar fasciitis [Mahowald et al], 5:385
Development of a novel rating system to assess lower-limb comfort [Kinchington et al], 5:371
Effect of foot orthoses contour on pain perception in individuals with patellofemoral pain [McPoil et al], 1:7
Patellofemoral pain syndrome and its association with hip, ankle, and foot function in 16- to 18-year-old high school students: a single-blind case-control study [Mølgaard et al], 3:215
Reducing the pain of local 1% lidocaine infiltration with a

preceding bacteriostatic saline injection: a double-blind prospective trial [Barrett et al], 3:223
Relationship between the size of bone marrow edema of the talus and ankle pain [Tonbul et al], 5:430
The symptomatic os vesalianum as an uncommon cause of lateral foot pain: a case report [Wilson et al], 4:356
Variations in foot posture and mobility between individuals with patellofemoral pain and those in a control group [McPoil et al], 4:289

Patella
Effect of foot orthoses contour on pain perception in individuals with patellofemoral pain [McPoil et al], 1:7
Patellofemoral pain syndrome and its association with hip, ankle, and foot function in 16- to 18-year-old high school students: a single-blind case-control study [Mølgaard et al], 3:215
Variations in foot posture and mobility between individuals with patellofemoral pain and those in a control group [McPoil et al], 4:289

Pediatric population
Patellofemoral pain syndrome and its association with hip, ankle, and foot function in 16- to 18-year-old high school students: a single-blind case-control study [Mølgaard et al], 3:215
Plantar pressures in children with and without Sever's disease [Becerro de Bengoa Vallejo et al], 1:17
Sever's disease: a prospective study of risk factors [Scharfbillig et al], 2:133
Small blue round cell tumor: an unusual case presentation in the foot [Devall et al], 4:363

Pedobarography
Surgical treatment of diaphyseal stress fractures of the fifth metatarsal in competitive athletes: long-term follow-up and computerized pedobarographic analysis [Smoljanovic et al], 6:517

Pes planus
Posterior tibial tendon dysfunction: a review [Durrant et al], 2:176

Posterior tibial tendon dysfunction
Posterior tibial tendon dysfunction: a review [Durrant et al], 2:176

Pregnancy
Changes in the plantar pressure distribution during gait throughout gestation [Ribeiro et al], 5:415

President's Message
Ambassador for the profession: the epilogue [Stone], 2:196
APMA House of Delegates: acceptance address [King], 3:278
Getting connected [King], 5:466
Giving back to podiatric medicine [King], 6:538
Planning for success [King], 4:370
The story of our success [Stone], 1:91

Pressure, plantar
Changes in the plantar pressure distribution during gait throughout gestation [Ribeiro et al], 5:415
Contralateral peak plantar pressures with a postoperative boot: a preliminary study [Mieras et al], 2:127
Evaluating intratester reliability of manual masking of plantar pressure measurements associated with chronic gout [Rome et al], 5:424
In-shoe plantar pressure distribution in nonneuropathic type 2 diabetic patients in Singapore [Tong et al], 6:509
Plantar pressures in children with and without Sever's disease [Becerro de Bengoa Vallejo et al], 1:17
Surgical treatment of pressure ulcers of the heel in skilled nursing facilities: a 12-year retrospective study of 57 patients [Han and Ezquerro], 2:167

Pronation
Foot orthoses: a review focusing on kinematics [Chevalier and Chockalingam], 4:341
The medial oblique shell inclination technique: a method to increase subtalar supination moments in foot orthoses [Harradine et al], 6:523
Patellofemoral pain syndrome and its association with hip, ankle, and foot function in 16- to 18-year-old high school students: a single-blind case-control study [Mølgaard et al], 3:215
Stabilization of joint forces of the subtalar complex via HyProCure sinus tarsi stent [Graham et al], 5:390
Variations in foot posture and mobility between individuals with patellofemoral pain and those in a control group [McPoil et al], 4:289

Pseudogout
Mimicry in older patients: tophaceous pseudogout as a tumorlike lesion: a case report [Erhardt et al], 5:462

R

Radiation
Invasive cutaneous squamous cell carcinoma presenting as a nonhealing radiation wound: a case report [Longobardi et al], 4:360

Reactant, acute phase
Acute phase reactants predict the risk of amputation in diabetic foot infection [Akinci et al], 1:1

Rheumatology
Evaluating intratester reliability of manual masking of plantar pressure measurements associated with chronic gout [Rome et al], 5:424
Reliability and validity of the INFOOT three-dimensional foot digitizer for patients with rheumatoid arthritis [De Mits et al], 3:198

Running
Barefoot running claims and controversies: a review of the literature [Jenkins and Cauthon], 3:231
Navicular stress reactions in runners: a review of evaluation and management of a competitive athlete [Yoho and Wells], 5:447

S

Saline
Reducing the pain of local 1% lidocaine infiltration with a preceding bacteriostatic saline injection: a double-blind prospective trial [Barrett et al], 3:223
The use of superoxidized aqueous solution versus saline as a replacement solution in the Versajet lavage system in chronic diabetic foot ulcers: a pilot study [Bowling et al], 2:124

Sarcoma
Small blue round cell tumor: an unusual case presentation in the foot [Devall et al], 4:363

Sesamoid
Bilateral tibial hallux sesamoid agenesis and fibular hallux sesamoid hypoplasia in a patient with bilateral hallux valgus [Marinozzi et al], 5:452

Sever's disease
Plantar pressures in children with and without Sever's disease [Becerro de Bengoa Vallejo et al], 1:17
Sever's disease: a prospective study of risk factors [Scharfbillig et al], 2:133

Silicone
A diabetic foot ulcer pilot study: does silicone gel sheeting reduce the incidence of reulceration? [Westphal et al], 2:116

Sinus tarsi
Stabilization of joint forces of the subtalar complex via HyProCure sinus tarsi stent [Graham et al], 5:390

Skin
Autologous skin fibroblast and keratinocyte grafts in the treatment of chronic foot ulcers in aging type 2 diabetic patients [Monami et al], 1:55
Clinical evaluation of 35% urea in a water-lipid–based foam containing lactic acid for treatment of mild-to-moderate xerosis of the foot [Grossman], 2:153
Invasive cutaneous squamous cell carcinoma presenting as a nonhealing radiation wound: a case report [Longobardi et al], 4:360
Prevalence of plantar verrucae in patients with human immunodeficiency virus infection during the post–highly active antiretroviral therapy era [Johnston et al], 1:35
Quantification of skin marker movement at the malleoli and talar heads [Birch and Deschamps], 6:497
Tungiasis in a beach volleyball player: a case report [Veraldi et al], 4:353

Splinting
Dynamic splinting for postoperative hallux limitus: a randomized, controlled trial [John et al], 4:285

Sports medicine
Barefoot running claims and controversies: a review of the literature [Jenkins and Cauthon], 3:231
Development of a novel rating system to assess lower-limb comfort [Kinchington et al], 5:371
Navicular stress reactions in runners: a review of evaluation and management of a competitive athlete [Yoho and Wells], 5:447
Surgical treatment of diaphyseal stress fractures of the fifth metatarsal in competitive athletes: long-term follow-up and computerized pedobarographic analysis [Smoljanovic et al], 6:517
Tungiasis in a beach volleyball player: a case report [Veraldi et al], 4:353

Stent
Stabilization of joint forces of the subtalar complex via HyProCure sinus tarsi stent [Graham et al], 5:390

Stress
Navicular stress reactions in runners: a review of evaluation and management of a competitive athlete [Yoho and Wells], 5:447
Surgical treatment of diaphyseal stress fractures of the fifth metatarsal in competitive athletes: long-term follow-up and computerized pedobarographic analysis [Smoljanovic et al], 6:517

Surgery
Central talar dome lesions: a unique surgical approach with incorporation of a talar allograft for joint reconstitution and restoration of function [Dobbs et al], 2:192
Distally based adipofascial sural flap for foot and ankle reconstruction [Mojallal et al], 1:41
Dorsal approach to transfer of the flexor digitorum brevis tendon [Becerro de Bengoa Vallejo et al], 4:297
Dynamic splinting for postoperative hallux limitus: a randomized, controlled trial [John et al], 4:285
Foot salvage after loss of the first and second metatarsal rays with a free fibular osteocutaneous flap [Salgado et al], 6:531
In vitro antimicrobial activity of calcium sulfate and hydroxyapatite (Cerament Bone Void Filler) discs using heat-sensitive and non–heat-sensitive antibiotics against methicillin-resistant *Staphylococcus aureus* and *Pseudomonas aeruginosa* [Karr et al], 2:146
Lengthening of the midfoot in a case of lower-extremity hemimelia [Kraus et al], 5:456
Management of grade III open dislocated ankle fractures: combined internal fixation with bioabsorbable screws/rods and external fixation [Ye et al], 4:307
Postoperative infection after excisional toenail matrixectomy: a retrospective clinical audit [Rusmir and Salerno], 4:316
Surgical treatment of diaphyseal stress fractures of the fifth metatarsal in competitive athletes: long-term follow-up and computerized pedobarographic analysis [Smoljanovic et al], 6:517
Surgical treatment of pressure ulcers of the heel in skilled nursing facilities: a 12-year retrospective study of 57 patients [Han and Ezquerro], 2:167
What do I do when bone hits the ground? [Brownell], 2:190

Synovitis
Giant cell tumor of the flexor hallucis longus tendon sheath: a case study [Findling et al], 2:187
Pigmented villonodular synovitis of the ankle: radiologic characteristics [Hao et al], 3:252

Synovium
Giant cell tumor of the flexor hallucis longus tendon sheath: a case study [Findling et al], 2:187
Pigmented villonodular synovitis of the ankle: radiologic characteristics [Hao et al], 3:252
Small blue round cell tumor: an unusual case presentation in the foot [Devall et al], 4:363

T

Talus
Central talar dome lesions: a unique surgical approach with incorporation of a talar allograft for joint reconstitution and restoration of function [Dobbs et al], 2:192
Quantification of skin marker movement at the malleoli and talar heads [Birch and Deschamps], 6:497
Relationship between the size of bone marrow edema of the talus and ankle pain [Tonbul et al], 5:430
An unusual cause of a cystic lesion with an osteochondral defect in the talus: intraosseous lipoma [Atay et al], 3:269
Validation of the talar–second metatarsal angle as a standard measurement for radiographic evaluation [Graham et al], 6:475

Tendon
Dorsal approach to transfer of the flexor digitorum brevis tendon [Becerro de Bengoa Vallejo et al], 4:297
Giant cell tumor of the flexor hallucis longus tendon sheath: a case study [Findling et al], 2:187
Posterior tibial tendon dysfunction: a review [Durrant et al], 2:176

Toe
Human digit partially consumed by a canine during sleep in a patient with neuropathy and diabetes [Rogers and Bevilacqua], 3:275
Postoperative infection after excisional toenail matrixectomy: a retrospective clinical audit [Rusmir and Salerno], 4:316
Prevalence of toenail onychomycosis in patients with type 2 diabetes mellitus and evaluation of risk factors [Gulcan et al], 1:49
A quantitative method for measuring forces applied by nail braces [Erdogan], 3:247

Tumor
Extra-abdominal desmoid fibromatosis in the foot: a case study [MacGill et al], 1:70
Giant cell tumor of the flexor hallucis longus tendon sheath: a case study [Findling et al], 2:187
Invasive cutaneous squamous cell carcinoma presenting as a nonhealing radiation wound: a case report [Longobardi et al], 4:360
Metastatic renal cell carcinoma presenting as foot metastasis: case report and review of the literature [Choufani et al], 3:265

Mimicry in older patients: tophaceous pseudogout as a tumorlike lesion: a case report [Erhardt et al], 5:462
Pigmented villonodular synovitis of the ankle: radiologic characteristics [Hao et al], 3:252
Small blue round cell tumor: an unusual case presentation in the foot [Devall et al], 4:363
An unusual cause of a cystic lesion with an osteochondral defect in the talus: intraosseous lipoma [Atay et al], 3:269

Tungiasis
Tungiasis in a beach volleyball player: a case report [Veraldi et al], 4:353

U

Ulcer
Acute phase reactants predict the risk of amputation in diabetic foot infection [Akinci et al], 1:1
Autologous skin fibroblast and keratinocyte grafts in the treatment of chronic foot ulcers in aging type 2 diabetic patients [Monami et al], 1:55
A diabetic foot ulcer pilot study: does silicone gel sheeting reduce the incidence of reulceration? [Westphal et al], 2:116
The economic value of specialized lower-extremity medical care by podiatric physicians in the treatment of diabetic foot ulcers [Carls et al], 2:93
Invasive cutaneous squamous cell carcinoma presenting as a nonhealing radiation wound: a case report [Longobardi et al], 4:360
Management in the wound-care center outpatient setting of a diabetic patient with forefoot osteomyelitis using Cerament Bone Void Filler impregnated with vancomycin: off-label use [Karr], 3:259
An open-label, three-arm pilot study of the safety and efficacy of topical Microcyn Rx wound care versus oral levofloxacin versus combined therapy for mild diabetic foot infections [Landsman et al], 6:484
Surgical treatment of pressure ulcers of the heel in skilled nursing facilities: a 12-year retrospective study of 57 patients [Han and Ezquerro], 2:167
The use of superoxidized aqueous solution versus saline as a replacement solution in the Versajet lavage system in chronic diabetic foot ulcers: a pilot study [Bowling et al], 2:124

Ultrasound
The correlation between plantar fascia thickness and symptoms of plantar fasciitis [Mahowald et al], 5:385

Urea
Clinical evaluation of 35% urea in a water-lipid–based foam containing lactic acid for treatment of mild-to-moderate xerosis of the foot [Grossman], 2:153

V

Vancomycin
Management in the wound-care center outpatient setting of a diabetic patient with forefoot osteomyelitis using Cerament Bone Void Filler impregnated with vancomycin: off-label use [Karr], 3:259

Verruca
Prevalence of plantar verrucae in patients with human immunodeficiency virus infection during the post–highly active antiretroviral therapy era [Johnston et al], 1:35

Versajet lavage system
The use of superoxidized aqueous solution versus saline as a replacement solution in the Versajet lavage system in chronic diabetic foot ulcers: a pilot study [Bowling et al], 2:124

Veterans
Prevalence of podiatric medical problems in veterans versus nonveterans [Shibuya et al], 4:323

Virus
Prevalence of plantar verrucae in patients with human immunodeficiency virus infection during the post–highly active antiretroviral therapy era [Johnston et al], 1:35

W

Walking
Changes in the plantar pressure distribution during gait throughout gestation [Ribeiro et al], 5:415
Evaluating intratester reliability of manual masking of plantar pressure measurements associated with chronic gout [Rome et al], 5:424
Foot orthoses: a review focusing on kinematics [Chevalier and Chockalingam], 4:341
In-shoe plantar pressure distribution in nonneuropathic type 2 diabetic patients in Singapore [Tong et al], 6:509
The in vitro reliability of the CODA MPX30 as the basis for a method of assessing the in vivo motion of the subtalar joint [Birch and Deschamps], 5:400
Quantification of skin marker movement at the malleoli and talar heads [Birch and Deschamps], 6:497
Soft-tissue movement at the foot during the stance phase of walking [Chen et al], 1:25

Wart
Prevalence of plantar verrucae in patients with human immunodeficiency virus infection during the post–highly active antiretroviral therapy era [Johnston et al], 1:35

X

Xerosis
Clinical evaluation of 35% urea in a water-lipid–based foam containing lactic acid for treatment of mild-to-moderate xerosis of the foot [Grossman], 2:153

AUTHOR INDEX

A

Acharya, U. Rajendra: In-shoe plantar pressure distribution in nonneuropathic type 2 diabetic patients in Singapore [Tong et al], 6:509

Akinci, Baris: Acute phase reactants predict the risk of amputation in diabetic foot infection [Akinci et al], 1:1

Akman, Budak: An unusual cause of a cystic lesion with an osteochondral defect in the talus: intraosseous lipoma [Atay et al], 3:269

American Diabetes Association: Foot care recommendations: an excerpt from *Diabetes Care*, 1:75

Aragón-Sánchez, Francisco Javier: Foot biomechanics in patients with diabetes mellitus: doubts regarding the relationship between neuropathy, foot motion, and deformities [Lázaro-Martínez et al], 3:208

Armstrong, David G.:

The Charcot foot in diabetes [Rogers et al], 5:437

Motivational interviewing by podiatric physicians: a method for improving patient self-care of the diabetic foot [Gabbay et al], 1:78

The use of superoxidized aqueous solution versus saline as a replacement solution in the Versajet lavage system in chronic diabetic foot ulcers: a pilot study [Bowling et al], 2:124

Atay, Evren Fehmi: An unusual cause of a cystic lesion with an osteochondral defect in the talus: intraosseous lipoma [Atay et al], 3:269

B

Baca, Emre: Relationship between the size of bone marrow edema of the talus and ankle pain [Tonbul et al], 5:430

Bagalman, J. Erin: The economic value of specialized lower-extremity medical care by podiatric physicians in the treatment of diabetic foot ulcers [Carls et al], 2:93

Ball, Kevin: Development of a novel rating system to assess lower-limb comfort [Kinchington et al], 5:371

Barbosa, Peter: Prevalence of plantar verrucae in patients with human immunodeficiency virus infection during the post–highly active antiretroviral therapy era [Johnston et al], 1:35

Barrett, Stephen L.:

Contralateral peak plantar pressures with a postoperative boot: a preliminary study [Mieras et al], 2:127

Reducing the pain of local 1% lidocaine infiltration with a preceding bacteriostatic saline injection: a double-blind prospective trial [Barrett et al], 3:223

Başsorgun, Cumhur Ibrahim: An unusual cause of a cystic lesion with an osteochondral defect in the talus: intraosseous lipoma [Atay et al], 3:269

Bayraktar, Firat: Acute phase reactants predict the risk of amputation in diabetic foot infection [Akinci et al], 1:1

Becerro de Bengoa Vallejo, Ricardo:

Dorsal approach to transfer of the flexor digitorum brevis tendon [Becerro de Bengoa Vallejo et al], 4:297

Plantar pressures in children with and without Sever's disease [Becerro de Bengoa Vallejo et al], 1:17

Beneit-Montesinos, Juan Vicente: Foot biomechanics in patients with diabetes mellitus: doubts regarding the relationship between neuropathy, foot motion, and deformities [Lázaro-Martínez et al], 3:208

Bes, Cemal: An unusual cause of a cystic lesion with an osteochondral defect in the talus: intraosseous lipoma [Atay et al], 3:269

Bevan, Lawrence: The medial oblique shell inclination technique: a method to increase subtalar supination moments in foot orthoses [Harradine et al], 6:523

Bevilacqua, Nicholas J.: Human digit partially consumed by a canine during sleep in a patient with neuropathy and diabetes [Rogers and Bevilacqua], 3:275

Birch, Ivan:

The in vitro reliability of the CODA MPX30 as the basis for a method of assessing the in vivo motion of the subtalar joint [Birch and Deschamps], 5:400

Quantification of skin marker movement at the malleoli and talar heads [Birch and Deschamps], 6:497

Blume, Peter A.: An open-label, three-arm pilot study of the safety and efficacy of topical Microcyn Rx wound care versus oral levofloxacin versus combined therapy for mild diabetic foot infections [Landsman et al], 6:484

Bojanic, Ivan: Surgical treatment of diaphyseal stress fractures of the fifth metatarsal in competitive athletes: long-term follow-up and computerized pedobarographic analysis [Smoljanovic et al], 6:517

Boulton, Andrew J.M.:

The Charcot foot in diabetes [Rogers et al], 5:437

The use of superoxidized aqueous solution versus saline as a replacement solution in the Versajet lavage system in chronic diabetic foot ulcers: a pilot study [Bowling et al], 2:124

Bower, Virginia M.: A diabetic foot ulcer pilot study: does silicone gel sheeting reduce the incidence of reulceration? [Westphal et al], 2:116

Bowling, Frank L.: The use of superoxidized aqueous solution versus saline as a replacement solution in the Versajet lavage system in chronic diabetic foot ulcers: a pilot study [Bowling et al], 2:124

Braye, Fabienne: Distally based adipofascial sural flap for foot and ankle reconstruction [Mojallal et al], 1:41

Breton, Pierre: Distally based adipofascial sural flap for foot and ankle reconstruction [Mojallal et al], 1:41

Brownell, Benjamin: What do I do when bone hits the ground? 2:190

C

Çakar, Murat: An unusual cause of a cystic lesion with an osteochondral defect in the talus: intraosseous lipoma [Atay et al], 3:269

Caldwell, Bryan D.: The use of focused electronic medical record forms to improve health-care outcomes [Caldwell et al], 4:331

Camison, Liliana: Foot salvage after loss of the first and second metatarsal rays with a free fibular osteocutaneous flap [Salgado et al], 6:531

Carls, Ginger S.: The economic value of specialized lower-extremity medical care by podiatric physicians in the treatment of diabetic foot ulcers [Carls et al], 2:93

Cauthon, David J.: Barefoot running claims and controversies: a review of the literature [Jenkins and Cauthon], 3:231

Cazzato, Luigi: Bilateral tibial hallux sesamoid agenesis and fibular hallux sesamoid hypoplasia in a patient with bilateral hallux valgus [Marinozzi et al], 5:452

Cazzell, Shawn M.: Central talar dome lesions: a unique surgical approach with incorporation of a talar allograft for joint reconstitution and restoration of function [Dobbs et al], 2:192

Chen, Aimin: Management of grade III open dislocated ankle fractures: combined internal fixation with bioabsorbable screws/rods and external fixation [Ye et al], 4:307

Chen, Shing-Jye: Soft-tissue movement at the foot during the stance phase of walking [Chen et al], 1:25

Chevalier, Thierry Larose: Foot orthoses: a review focusing on kinematics [Chevalier and Chockalingam], 4:341

Chikka, Avanthi: Validation of the talar–second metatarsal angle as a standard measurement for radiographic evaluation [Graham et al], 6:475

Chockalingam, Nachiappan:

Clinical assessment of ankle joint dorsiflexion: a review of measurement techniques [Gatt and Chockalingam], 1:59

Foot orthoses: a review focusing on kinematics [Chevalier and Chockalingam], 4:341
Posterior tibial tendon dysfunction: a review [Durrant et al], 2:176
Chou, Li-Shan: Soft-tissue movement at the foot during the stance phase of walking [Chen et al], 1:25
Choufani, Elie: Metastatic renal cell carcinoma presenting as foot metastasis: case report and review of the literature [Choufani et al], 3:265
Christina, James R.: The economic value of specialized lower-extremity medical care by podiatric physicians in the treatment of diabetic foot ulcers [Carls et al], 2:93
Chua, Kuang C.: In-shoe plantar pressure distribution in nonneuropathic type 2 diabetic patients in Singapore [Tong et al], 6:509
Chun, Kyung Ah: Freiberg's disease: quantitative assessment of osteonecrosis on three-dimensional CT [Chun et al], 4:335
Ciliberti, Louis J., Jr.: Prevalence of podiatric medical problems in veterans versus nonveterans [Shibuya et al], 4:323
Collins, Simon: The medial oblique shell inclination technique: a method to increase subtalar supination moments in foot orthoses [Harradine et al], 6:523
Cooper, Kimbal E.: Reducing the pain of local 1% lidocaine infiltration with a preceding bacteriostatic saline injection: a double-blind prospective trial [Barrett et al], 3:223
Coorevits, Pascal: Reliability and validity of the INFOOT three-dimensional foot digitizer for patients with rheumatoid arthritis [De Mits et al], 3:198
Cornwall, Mark W.:
Effect of foot orthoses contour on pain perception in individuals with patellofemoral pain [McPoil et al], 1:7
Variations in foot posture and mobility between individuals with patellofemoral pain and those in a control group [McPoil et al], 4:289
Crews, Ryan T.: The use of superoxidized aqueous solution versus saline as a replacement solution in the Versajet lavage system in chronic diabetic foot ulcers: a pilot study [Bowling et al], 2:124

D

Dalbeth, Nicola: Evaluating intratester reliability of manual masking of plantar pressure measurements associated with chronic gout [Rome et al], 5:424
Davenport, Colin: Charcot-Marie-Tooth disease complicating type 2 diabetes [Win et al], 4:349
De Clercq, Dirk: Reliability and validity of the INFOOT three-dimensional foot digitizer for patients with rheumatoid arthritis [De Mits et al], 3:198
DeFrancis, Roy R.: The economic value of specialized lower-extremity medical care by podiatric physicians in the treatment of diabetic foot ulcers [Carls et al], 2:93
Delaney, Siobhan: Charcot-Marie-Tooth disease complicating type 2 diabetes [Win et al], 4:349
De Mits, Sophie: Reliability and validity of the INFOOT three-dimensional foot digitizer for patients with rheumatoid arthritis [De Mits et al], 3:198
Denaro, Vincenzo: Bilateral tibial hallux sesamoid agenesis and fibular hallux sesamoid hypoplasia in a patient with bilateral hallux valgus [Marinozzi et al], 5:452
Deschamps, Kevin:
The in vitro reliability of the CODA MPX30 as the basis for a method of assessing the in vivo motion of the subtalar joint [Birch and Deschamps], 5:400
Quantification of skin marker movement at the malleoli and talar heads [Birch and Deschamps], 6:497
Desideri, Carla Maria: Autologous skin fibroblast and keratinocyte grafts in the treatment of chronic foot ulcers in aging type 2 diabetic patients [Monami et al], 1:55
Devall, J. Marshall: Small blue round cell tumor: an unusual case presentation in the foot [Devall et al], 4:363
Diligent, Jerome: Metastatic renal cell carcinoma presenting as foot metastasis: case report and review of the literature [Choufani et al], 3:265
Dini, Monara: Central talar dome lesions: a unique surgical approach with incorporation of a talar allograft for joint reconstitution and restoration of function [Dobbs et al], 2:192
Dobbs, Bruce M.: Central talar dome lesions: a unique surgical approach with incorporation of a talar allograft for joint reconstitution and restoration of function [Dobbs et al], 2:192
Dominguez, Gabriel: Hallux limitus and its relationship with the internal rotational pattern of the lower limb [Munuera et al], 6:467
Driver, Vickie R.: The economic value of specialized lower-extremity medical care by podiatric physicians in the treatment of diabetic foot ulcers [Carls et al], 2:93
Duncan, Alissa N.: Foot salvage after loss of the first and second metatarsal rays with a free fibular osteocutaneous flap [Salgado et al], 6:531
Durrant, Beverley: Posterior tibial tendon dysfunction: a review [Durrant et al], 2:176

E

Edmonds, Michael: The Charcot foot in diabetes [Rogers et al], 5:437
Elewaut, Dirk: Reliability and validity of the INFOOT three-dimensional foot digitizer for patients with rheumatoid arthritis [De Mits et al], 3:198
Erdogan, Fatma G.: A quantitative method for measuring forces applied by nail braces, 3:247
Erhardt, Matthias A.: Mimicry in older patients: tophaceous pseudogout as a tumorlike lesion: a case report [Erhardt et al], 5:462
Ezquerro, Ruben: Surgical treatment of pressure ulcers of the heel in skilled nursing facilities: a 12-year retrospective study of 57 patients [Han and Ezquerro], 2:167

F

Findling, Jeff: Giant cell tumor of the flexor hallucis longus tendon sheath: a case study [Findling et al], 2:187
Frank, Christian B.: Mimicry in older patients: tophaceous pseudogout as a tumorlike lesion: a case report [Erhardt et al], 5:462
Freeman, Denise: Interrater and intrarater reliability in the measurement of ankle joint dorsiflexion is independent of examiner experience and technique used [Kim et al], 5:407
Frush, Katherine M.: Small blue round cell tumor: an unusual case presentation in the foot [Devall et al], 4:363
Frykberg, Robert G.: The Charcot foot in diabetes [Rogers et al], 5:437
Fuller, David A.: Foot salvage after loss of the first and second metatarsal rays with a free fibular osteocutaneous flap [Salgado et al], 6:531

G

Gabbay, Robert A.: Motivational interviewing by podiatric physicians: a method for improving patient self-care of the diabetic foot [Gabbay et al], 1:78
Galois, Laurent: Metastatic renal cell carcinoma presenting as foot metastasis: case report and review of the literature [Choufani et al], 3:265
Game, Frances: The Charcot foot in diabetes [Rogers et al], 5:437
García Morales, Esther: Foot biomechanics in patients with diabetes mellitus: doubts regarding the relationship between neuropathy, foot motion, and deformities [Lázaro-Martínez et al], 3:208
Garoufalis, Matthew G.: The economic value of specialized lower-extremity medical care by podiatric physicians in the treatment of diabetic foot ulcers [Carls et al], 2:93

Gatt, Alfred: Clinical assessment of ankle joint dorsiflexion: a review of measurement techniques [Gatt and Chockalingam], 1:59
Gibson, Teresa B.: The economic value of specialized lower-extremity medical care by podiatric physicians in the treatment of diabetic foot ulcers [Carls et al], 2:93
Goel, Vijay: Stabilization of joint forces of the subtalar complex via HyProCure sinus tarsi stent [Graham et al], 5:390
Gonen, Aysun: Relationship between the size of bone marrow edema of the talus and ankle pain [Tonbul et al], 5:430
Gonzáles-Jurado, Maximo A.: Foot biomechanics in patients with diabetes mellitus: doubts regarding the relationship between neuropathy, foot motion, and deformities [Lázaro-Martínez et al], 3:208
Gou, Sanhuai: Management of grade III open dislocated ankle fractures: combined internal fixation with bioabsorbable screws/rods and external fixation [Ye et al], 4:307
Grady, John F.: The correlation between plantar fascia thickness and symptoms of plantar fasciitis [Mahowald et al], 5:385
Graham, Michael E.:
Stabilization of joint forces of the subtalar complex via HyProCure sinus tarsi stent [Graham et al], 5:390
Validation of the talar–second metatarsal angle as a standard measurement for radiographic evaluation [Graham et al], 6:475
Groner, Thomas W.: Giant cell tumor of the flexor hallucis longus tendon sheath: a case study [Findling et al], 2:187
Grossman, Allan B.: Clinical evaluation of 35% urea in a water-lipid–based foam containing lactic acid for treatment of mild-to-moderate xerosis of the foot, 2:153
Gulcan, Aynur: Prevalence of toenail onychomycosis in patients with type 2 diabetes mellitus and evaluation of risk factors [Gulcan et al], 1:49
Gulcan, Erim: Prevalence of toenail onychomycosis in patients with type 2 diabetes mellitus and evaluation of risk factors [Gulcan et al], 1:49
Gurr, Joel M.: A diabetic foot ulcer pilot study: does silicone gel sheeting reduce the incidence of reulceration? [Westphal et al], 2:116
Gutierrez, Andres: An open-label, three-arm pilot study of the safety and efficacy of topical Microcyn Rx wound care versus oral levofloxacin versus combined therapy for mild diabetic foot infections [Landsman et al], 6:484
Güven, Melih: An unusual cause of a cystic lesion with an osteochondral defect in the talus: intraosseous lipoma [Atay et al], 3:269
Guzelant, Aliye Yildirim: Relationship between the size of bone marrow edema of the talus and ankle pain [Tonbul et al], 5:430

H

Han, Paul Y.: Surgical treatment of pressure ulcers of the heel in skilled nursing facilities: a 12-year retrospective study of 57 patients [Han and Ezquerro], 2:167
Hao, Da-Peng: Pigmented villonodular synovitis of the ankle: radiologic characteristics [Hao et al], 3:252
Harradine, Paul: The medial oblique shell inclination technique: a method to increase subtalar supination moments in foot orthoses [Harradine et al], 6:523
Harrison, Jessica C.: A diabetic foot ulcer pilot study: does silicone gel sheeting reduce the incidence of reulceration? [Westphal et al], 2:116
Hartemann, Agnes: The Charcot foot in diabetes [Rogers et al], 5:437
Hashmi, Farina: Posterior tibial tendon dysfunction: a review [Durrant et al], 2:176

I

Ippolito, Giulio: Autologous skin fibroblast and keratinocyte grafts in the treatment of chronic foot ulcers in aging type 2 diabetic patients [Monami et al], 1:55
Ivkovic, Alan: Surgical treatment of diaphyseal stress fractures of the fifth metatarsal in competitive athletes: long-term follow-up and computerized pedobarographic analysis [Smoljanovic et al], 6:517

J

Jeffcoate, William: The Charcot foot in diabetes [Rogers et al], 5:437
Jelic, Miroslav: Surgical treatment of diaphyseal stress fractures of the fifth metatarsal in competitive athletes: long-term follow-up and computerized pedobarographic analysis [Smoljanovic et al], 6:517
Jenkins, David W.: Barefoot running claims and controversies: a review of the literature [Jenkins and Cauthon], 3:231
Jirkovska, Alexandra: The Charcot foot in diabetes [Rogers et al], 5:437
João, Sílvia Maria Amado: Changes in the plantar pressure distribution during gait throughout gestation [Ribeiro et al], 5:415
John, Mathew M.: Dynamic splinting for postoperative hallux limitus: a randomized, controlled trial [John et al], 4:285
Johnston, James: Prevalence of plantar verrucae in patients with human immunodeficiency virus infection during the post–highly active antiretroviral therapy era [Johnston et al], 1:35
Jones, Paul C.: Validation of the talar–second metatarsal angle as a standard measurement for radiographic evaluation [Graham et al], 6:475
Jones, Sara: Sever's disease: a prospective study of risk factors [Scharfbillig et al], 2:133
Jordon, Douglas A., Jr.: An open-label, three-arm pilot study of the safety and efficacy of topical Microcyn Rx wound care versus oral levofloxacin versus combined therapy for mild diabetic foot infections [Landsman et al], 6:484
Joseph, Warren S.: Guidelines for authors, 1:85, 4:371
Jude, Edward: The Charcot foot in diabetes [Rogers et al], 5:437
Jules, Kevin T.: Dorsal approach to transfer of the flexor digitorum brevis tendon [Becerro de Bengoa Vallejo et al], 4:297
Jupiter, Daniel C.: Prevalence of podiatric medical problems in veterans versus nonveterans [Shibuya et al], 4:323

K

Kalish, Stanley: Dynamic splinting for postoperative hallux limitus: a randomized, controlled trial [John et al], 4:285
Karr, Jeffrey C.:
In vitro antimicrobial activity of calcium sulfate and hydroxyapatite (Cerament Bone Void Filler) discs using heat-sensitive and non–heat-sensitive antibiotics against methicillin-resistant *Staphylococcus aureus* and *Pseudomonas aeruginosa* [Karr et al], 2:146
Management in the wound-care center outpatient setting of a diabetic patient with forefoot osteomyelitis using Cerament Bone Void Filler impregnated with vancomycin: off-label use, 3:259
Katz, Robert D.: The use of focused electronic medical record forms to improve health-care outcomes [Caldwell et al], 4:331
Kaul, Shailja: Motivational interviewing by podiatric physicians: a method for improving patient self-care of the diabetic foot [Gabbay et al], 1:78
Kaya, Demet: Prevalence of toenail onychomycosis in patients with type 2 diabetes mellitus and evaluation of risk factors [Gulcan et al], 1:49
Kelly, Margaret: Charcot-Marie-Tooth disease complicating type 2 diabetes [Win et al], 4:349
Keriazes, Georgia: In vitro antimicrobial activity of calcium sulfate and hydroxyapatite (Cerament Bone Void Filler) discs using heat-sensitive and non–heat-sensitive antibiotics against

methicillin-resistant *Staphylococcus aureus* and *Pseudomonas aeruginosa* [Karr et al], 2:146
Khademi, Saieh: Prevalence of plantar verrucae in patients with human immunodeficiency virus infection during the post–highly active antiretroviral therapy era [Johnston et al], 1:35
Kim, Paul Jeong: Interrater and intrarater reliability in the measurement of ankle joint dorsiflexion is independent of examiner experience and technique used [Kim et al], 5:407
Kinchington, Michael: Development of a novel rating system to assess lower-limb comfort [Kinchington et al], 5:371
King, Christy M.: Prevalence of plantar verrucae in patients with human immunodeficiency virus infection during the post–highly active antiretroviral therapy era [Johnston et al], 1:35
King, Michael J.:
APMA House of Delegates: acceptance address, 3:278
Getting connected, 5:466
Giving back to podiatric medicine, 6:538
Planning for success, 4:370
Kraus, Tanja: Lengthening of the midfoot in a case of lower-extremity hemimelia [Kraus et al], 5:456
Kucukyavas, Yasin: Acute phase reactants predict the risk of amputation in diabetic foot infection [Akinci et al], 1:1

L

La Fontaine, Javier: Prevalence of podiatric medical problems in veterans versus nonveterans [Shibuya et al], 4:323
Lafuente, Blanca: Hallux limitus and its relationship with the internal rotational pattern of the lower limb [Munuera et al], 6:467
Lafuente, Guillermo: Hallux limitus and its relationship with the internal rotational pattern of the lower limb [Munuera et al], 6:467
Landsman, Adam: An open-label, three-arm pilot study of the safety and efficacy of topical Microcyn Rx wound care versus oral levofloxacin versus combined therapy for mild diabetic foot infections [Landsman et al], 6:484
LaScola, Natalie K.: Giant cell tumor of the flexor hallucis longus tendon sheath: a case study [Findling et al], 2:187
Lauretta, Joseph: In vitro antimicrobial activity of calcium sulfate and hydroxyapatite (Cerament Bone Void Filler) discs using heat-sensitive and non–heat-sensitive antibiotics against methicillin-resistant *Staphylococcus aureus* and *Pseudomonas aeruginosa* [Karr et al], 2:146
Lázaro-Martínez, José Luis: Foot biomechanics in patients with diabetes mellitus: doubts regarding the relationship between neuropathy, foot motion, and deformities [Lázaro-Martínez et al], 3:208
Legge, Bradford S.: The correlation between plantar fascia thickness and symptoms of plantar fasciitis [Mahowald et al], 5:385
Leino-Kilpi, Helena: Effect of an educational intervention on nurses' knowledge of foot care and on the foot health of older residents [Stolt et al], 2:159
Lin, Chih-Hung: Foot salvage after loss of the first and second metatarsal rays with a free fibular osteocutaneous flap [Salgado et al], 6:531
Linhart, Wolfgang E.: Lengthening of the midfoot in a case of lower-extremity hemimelia [Kraus et al], 5:456
Lobo, Maria: Evaluating intratester reliability of manual masking of plantar pressure measurements associated with chronic gout [Rome et al], 5:424
Longo, Umile Giuseppe: Bilateral tibial hallux sesamoid agenesis and fibular hallux sesamoid hypoplasia in a patient with bilateral hallux valgus [Marinozzi et al], 5:452
Longobardi, Stephen J.: Invasive cutaneous squamous cell carcinoma presenting as a nonhealing radiation wound: a case report [Longobardi et al], 4:360
López Chicharro, José: Plantar pressures in children with and without Sever's disease [Becerro de Bengoa Vallejo et al], 1:17
Losa Iglesias, Marta Elena:
Dorsal approach to transfer of the flexor digitorum brevis tendon [Becerro de Bengoa Vallejo et al], 4:297
Plantar pressures in children with and without Sever's disease [Becerro de Bengoa Vallejo et al], 1:17

M

MacGill, Alan A.: Extra-abdominal desmoid fibromatosis in the foot: a case study [MacGill et al], 1:70
Maffulli, Nicola: Bilateral tibial hallux sesamoid agenesis and fibular hallux sesamoid hypoplasia in a patient with bilateral hallux valgus [Marinozzi et al], 5:452
Mahowald, Sarah: The correlation between plantar fascia thickness and symptoms of plantar fasciitis [Mahowald et al], 5:385
Mainard, Didier: Metastatic renal cell carcinoma presenting as foot metastasis: case report and review of the literature [Choufani et al], 3:265
Mannucci, Edoardo: Autologous skin fibroblast and keratinocyte grafts in the treatment of chronic foot ulcers in aging type 2 diabetic patients [Monami et al], 1:55
Mansour, E. Hani: Invasive cutaneous squamous cell carcinoma presenting as a nonhealing radiation wound: a case report [Longobardi et al], 4:360
Marchionni, Niccolò: Autologous skin fibroblast and keratinocyte grafts in the treatment of chronic foot ulcers in aging type 2 diabetic patients [Monami et al], 1:55
Mardini, Samir: Foot salvage after loss of the first and second metatarsal rays with a free fibular osteocutaneous flap [Salgado et al], 6:531
Marinozzi, Andrea: Bilateral tibial hallux sesamoid agenesis and fibular hallux sesamoid hypoplasia in a patient with bilateral hallux valgus [Marinozzi et al], 5:452
Martinelli, Nicolò: Bilateral tibial hallux sesamoid agenesis and fibular hallux sesamoid hypoplasia in a patient with bilateral hallux valgus [Marinozzi et al], 5:452
Martínez Hernández, David: Foot biomechanics in patients with diabetes mellitus: doubts regarding the relationship between neuropathy, foot motion, and deformities [Lázaro-Martínez et al], 3:208
Matyas, Aaron: Stabilization of joint forces of the subtalar complex via HyProCure sinus tarsi stent [Graham et al], 5:390
Maxka, Jim: Reducing the pain of local 1% lidocaine infiltration with a preceding bacteriostatic saline injection: a double-blind prospective trial [Barrett et al], 3:223
McNair, Peter J.: Evaluating intratester reliability of manual masking of plantar pressure measurements associated with chronic gout [Rome et al], 5:424
McPoil, Thomas G.:
Effect of foot orthoses contour on pain perception in individuals with patellofemoral pain [McPoil et al], 1:7
Variations in foot posture and mobility between individuals with patellofemoral pain and those in a control group [McPoil et al], 4:289
McQueen, Fiona: Evaluating intratester reliability of manual masking of plantar pressure measurements associated with chronic gout [Rome et al], 5:424
Mhatre, Devdatt: Stabilization of joint forces of the subtalar complex via HyProCure sinus tarsi stent [Graham et al], 5:390
Mieras, Jamie N.:
Contralateral peak plantar pressures with a postoperative boot: a preliminary study [Mieras et al], 2:127
Interrater and intrarater reliability in the measurement of ankle joint dorsiflexion is independent of examiner experience and technique used [Kim et al], 5:407
Reducing the pain of local 1% lidocaine infiltration with a preceding bacteriostatic saline injection: a double-blind prospective trial [Barrett et al], 3:223
Millione, Vincent R.: Extra-abdominal desmoid fibromatosis in the foot: a case study [MacGill et al], 1:70
Mirkovic, Maja: Surgical treatment of diaphyseal stress fractures of the fifth metatarsal in competitive athletes: long-term

follow-up and computerized pedobarographic analysis [Smoljanovic et al], 6:517
Mojallal, Ali: Distally based adipofascial sural flap for foot and ankle reconstruction [Mojallal et al], 1:41
Mølgaard, Carsten: Patellofemoral pain syndrome and its association with hip, ankle, and foot function in 16- to 18-year-old high school students: a single-blind case-control study [Mølgaard et al], 3:517
Monami, Matteo: Autologous skin fibroblast and keratinocyte grafts in the treatment of chronic foot ulcers in aging type 2 diabetic patients [Monami et al], 1:55
Morbach, Stephan: The Charcot foot in diabetes [Rogers et al], 5:437
Morrison, William B.: The Charcot foot in diabetes [Rogers et al], 5:437
Mukul, Mukherjee: Soft-tissue movement at the foot during the stance phase of walking [Chen et al], 1:25
Munuera, Pedro V.: Hallux limitus and its relationship with the internal rotational pattern of the lower limb [Munuera et al], 6:467

N

Narayanswamy, C.: Anatomical variations of peroneal muscles: a cadaver study in an Indian population and a review of the literature [Prakash et al], 6:505
Naughton, Geraldine: Development of a novel rating system to assess lower-limb comfort [Kinchington et al], 5:371
Neame, Ingrid M.: A diabetic foot ulcer pilot study: does silicone gel sheeting reduce the incidence of reulceration? [Westphal et al], 2:116
Nelson, Joseph: Prevalence of plantar verrucae in patients with human immunodeficiency virus infection during the post–highly active antiretroviral therapy era [Johnston et al], 1:35

O

Oh, Hyung Keun: Freiberg's disease: quantitative assessment of osteonecrosis on three-dimensional CT [Chun et al], 4:335
Oksuz, Sukru: Prevalence of toenail onychomycosis in patients with type 2 diabetes mellitus and evaluation of risk factors [Gulcan et al], 1:49
Ouzounov, Kaloian: The symptomatic os vesalianum as an uncommon cause of lateral foot pain: a case report [Wilson et al], 4:356
Ozbaydar, Mehmet U.: Relationship between the size of bone marrow edema of the talus and ankle pain [Tonbul et al], 5:430

P

Page, Jeffrey: Interrater and intrarater reliability in the measurement of ankle joint dorsiflexion is independent of examiner experience and technique used [Kim et al], 5:407
Parikh, Rachit: Stabilization of joint forces of the subtalar complex via HyProCure sinus tarsi stent [Graham et al], 5:390
Pascarella, Eugene M.: The use of focused electronic medical record forms to improve health-care outcomes [Caldwell et al], 4:331
Peace, Ruth: Interrater and intrarater reliability in the measurement of ankle joint dorsiflexion is independent of examiner experience and technique used [Kim et al], 5:407
Pecina, Marko: Surgical treatment of diaphyseal stress fractures of the fifth metatarsal in competitive athletes: long-term follow-up and computerized pedobarographic analysis [Smoljanovic et al], 6:517
Perns, Stephen V.: Dynamic splinting for postoperative hallux limitus: a randomized, controlled trial [John et al], 4:285
Persico, Maria Chiara: Tungiasis in a beach volleyball player: a case report [Veraldi et al], 4:353
Pinzur, Michael: The Charcot foot in diabetes [Rogers et al], 5:437
Pitocco, Dario: The Charcot foot in diabetes [Rogers et al], 5:437
Prados Frutos, Carlos:
Dorsal approach to transfer of the flexor digitorum brevis tendon [Becerro de Bengoa Vallejo et al], 4:297
Plantar pressures in children with and without Sever's disease [Becerro de Bengoa Vallejo et al], 1:17
Prakash: Anatomical variations of peroneal muscles: a cadaver study in an Indian population and a review of the literature [Prakash et al], 6:505

R

Rajini, Thimmiah: Anatomical variations of peroneal muscles: a cadaver study in an Indian population and a review of the literature [Prakash et al], 6:505
Rathleff, Michael Skovdal: Patellofemoral pain syndrome and its association with hip, ankle, and foot function in 16- to 18-year-old high school students: a single-blind case-control study [Mølgaard et al], 3:215
Reina, Maria: Hallux limitus and its relationship with the internal rotational pattern of the lower limb [Munuera et al], 6:467
Ribeiro, Ana Paula: Changes in the plantar pressure distribution during gait throughout gestation [Ribeiro et al], 5:415
Rodríguez, Miguel Fuentes: Dorsal approach to transfer of the flexor digitorum brevis tendon [Becerro de Bengoa Vallejo et al], 4:297
Rodríguez Sanz, David: Plantar pressures in children with and without Sever's disease [Becerro de Bengoa Vallejo et al], 1:17
Rogers, Lee C.:
The Charcot foot in diabetes [Rogers et al], 5:437
Human digit partially consumed by a canine during sleep in a patient with neuropathy and diabetes [Rogers and Bevilacqua], 3:275
Rome, Keith: Evaluating intratester reliability of manual masking of plantar pressure measurements associated with chronic gout [Rome et al], 5:424
Roosen, Philip: Reliability and validity of the INFOOT three-dimensional foot digitizer for patients with rheumatoid arthritis [De Mits et al], 3:198
Routasalo, Pirkko: Effect of an educational intervention on nurses' knowledge of foot care and on the foot health of older residents [Stolt et al], 2:159
Ruano, Rodrigo: Changes in the plantar pressure distribution during gait throughout gestation [Ribeiro et al], 5:415
Rusmir, Alen: Postoperative infection after excisional toenail matrixectomy: a retrospective clinical audit [Rusmir and Salerno], 4:316

S

Sacco, Isabel de Camargo Neves: Changes in the plantar pressure distribution during gait throughout gestation [Ribeiro et al], 5:415
Sahin, Idris: Prevalence of toenail onychomycosis in patients with type 2 diabetes mellitus and evaluation of risk factors [Gulcan et al], 1:49
Salerno, Angelo: Postoperative infection after excisional toenail matrixectomy: a retrospective clinical audit [Rusmir and Salerno], 4:316
Salgado, Christopher J.: Foot salvage after loss of the first and second metatarsal rays with a free fibular osteocutaneous flap [Salgado et al], 6:531
Salgami, Eleanna: The use of superoxidized aqueous solution versus saline as a replacement solution in the Versajet lavage system in chronic diabetic foot ulcers: a pilot study [Bowling et al], 2:124
Salvadores Fuentes, Paloma: Plantar pressures in children with and without Sever's disease [Becerro de Bengoa Vallejo et al], 1:17
Sanders, Lee: The Charcot foot in diabetes [Rogers et al], 5:437

Scharfbillig, Rolf W.: Sever's disease: a prospective study of risk factors [Scharfbillig et al], 2:133
Scheffler, Neil M.: Motivational interviewing by podiatric physicians: a method for improving patient self-care of the diabetic foot [Gabbay et al], 1:78
Scutter, Sheila: Sever's disease: a prospective study of risk factors [Scharfbillig et al], 2:133
Shanks, Sky: Prevalence of plantar verrucae in patients with human immunodeficiency virus infection during the post–highly active antiretroviral therapy era [Johnston et al], 1:35
Shibuya, Naohiro: Prevalence of podiatric medical problems in veterans versus nonveterans [Shibuya et al], 4:323
Shipkov, Christo D.: Distally based adipofascial sural flap for foot and ankle reconstruction [Mojallal et al], 1:41
Simonsen, Ole: Patellofemoral pain syndrome and its association with hip, ankle, and foot function in 16- to 18-year-old high school students: a single-blind case-control study [Mølgaard et al], 3:215
Singh, Deepak Kumar: Anatomical variations of peroneal muscles: a cadaver study in an Indian population and a review of the literature [Prakash et al], 6:505
Singh, Gajendra: Anatomical variations of peroneal muscles: a cadaver study in an Indian population and a review of the literature [Prakash et al], 6:505
Singleton, Tanya J.: Contralateral peak plantar pressures with a postoperative boot: a preliminary study [Mieras et al], 2:127
Smith, Diarmuid: Charcot-Marie-Tooth disease complicating type 2 diabetes [Win et al], 4:349
Smoljanovic, Tomislav: Surgical treatment of diaphyseal stress fractures of the fifth metatarsal in competitive athletes: long-term follow-up and computerized pedobarographic analysis [Smoljanovic et al], 6:517
Steiner, Lori: Small blue round cell tumor: an unusual case presentation in the foot [Devall et al], 4:363
Stolt, Minna: Effect of an educational intervention on nurses' knowledge of foot care and on the foot health of older residents [Stolt et al], 2:159
Stone, Kathleen M.:
Ambassador for the profession: the epilogue, 2:196
The story of our success, 1:91
Suda, Arnold J.: Mimicry in older patients: tophaceous pseudogout as a tumorlike lesion: a case report [Erhardt et al], 5:462
Suh, Jin Soo: Freiberg's disease: quantitative assessment of osteonecrosis on three-dimensional CT [Chun et al], 4:335
Suhonen, Riitta: Effect of an educational intervention on nurses' knowledge of foot care and on the foot health of older residents [Stolt et al], 2:159
Sullivan, Brian: Invasive cutaneous squamous cell carcinoma presenting as a nonhealing radiation wound: a case report [Longobardi et al], 4:360
Sullivan, Laura G.: Extra-abdominal desmoid fibromatosis in the foot: a case study [MacGill et al], 1:70
Survepalli, David George: Evaluating intratester reliability of manual masking of plantar pressure measurements associated with chronic gout [Rome et al], 5:424
Svehlik, Martin: Lengthening of the midfoot in a case of lower-extremity hemimelia [Kraus et al], 5:456

T

Tan, Peck H.: In-shoe plantar pressure distribution in non-neuropathic type 2 diabetic patients in Singapore [Tong et al], 6:509
Thoms, Tanya: Interrater and intrarater reliability in the measurement of ankle joint dorsiflexion is independent of examiner experience and technique used [Kim et al], 5:407
Tonbul, Murat: Relationship between the size of bone marrow edema of the talus and ankle pain [Tonbul et al], 5:430
Tong, Jasper W.K.: In-shoe plantar pressure distribution in nonneuropathic type 2 diabetic patients in Singapore [Tong et al], 6:509
Trombini-Souza, Francis: Changes in the plantar pressure distribution during gait throughout gestation [Ribeiro et al], 5:415

U

Uccioli, Luigi: The Charcot foot in diabetes [Rogers et al], 5:437
Ulbrecht, Jan: Motivational interviewing by podiatric physicians: a method for improving patient self-care of the diabetic foot [Gabbay et al], 1:78

V

Valsecchi, Marta: Tungiasis in a beach volleyball player: a case report [Veraldi et al], 4:353
Van, Georges Ha: The Charcot foot in diabetes [Rogers et al], 5:437
VanBuren, Vincent: Prevalence of podiatric medical problems in veterans versus nonveterans [Shibuya et al], 4:323
Vayser, Dean: An open-label, three-arm pilot study of the safety and efficacy of topical Microcyn Rx wound care versus oral levofloxacin versus combined therapy for mild diabetic foot infections [Landsman et al], 6:484
Venkatiah, Jayanthi: Anatomical variations of peroneal muscles: a cadaver study in an Indian population and a review of the literature [Prakash et al], 6:505
Veraldi, Stefano: Tungiasis in a beach volleyball player: a case report [Veraldi et al], 4:353
Vetter, Sven Y.: Mimicry in older patients: tophaceous pseudogout as a tumorlike lesion: a case report [Erhardt et al], 5:462
Vicenzino, Bill:
Effect of foot orthoses contour on pain perception in individuals with patellofemoral pain [McPoil et al], 1:7
Variations in foot posture and mobility between individuals with patellofemoral pain and those in a control group [McPoil et al], 4:289
Vivarelli, Maria: Autologous skin fibroblast and keratinocyte grafts in the treatment of chronic foot ulcers in aging type 2 diabetic patients [Monami et al], 1:55

W

Wang, Kook Hyun: Freiberg's disease: quantitative assessment of osteonecrosis on three-dimensional CT [Chun et al], 4:335
Wang, Shaohung: The economic value of specialized lower-extremity medical care by podiatric physicians in the treatment of diabetic foot ulcers [Carls et al], 2:93
Wang, Xue-Na: Pigmented villonodular synovitis of the ankle: radiologic characteristics [Hao et al], 3:252
Wang, Zhen-Chang: Pigmented villonodular synovitis of the ankle: radiologic characteristics [Hao et al], 3:252
Warren, Meghan: Variations in foot posture and mobility between individuals with patellofemoral pain and those in a control group [McPoil et al], 4:289
Webb, Chris: The medial oblique shell inclination technique: a method to increase subtalar supination moments in foot orthoses [Harradine et al], 6:523
Wells, Shevonne K.: Navicular stress reactions in runners: a review of evaluation and management of a competitive athlete [Yoho and Wells], 5:447
Wentzensen, Andreas: Mimicry in older patients: tophaceous pseudogout as a tumorlike lesion: a case report [Erhardt et al], 5:462
Westphal, Cara: A diabetic foot ulcer pilot study: does silicone gel sheeting reduce the incidence of reulceration? [Westphal et al], 2:116
Willis, F. Buck: Dynamic splinting for postoperative hallux limitus: a randomized, controlled trial [John et al], 4:285
Wilson, Richard C.: The symptomatic os vesalianum as an uncommon cause of lateral foot pain: a case report [Wilson et al], 4:356

Wilson, Thomas C.: The symptomatic os vesalianum as an uncommon cause of lateral foot pain: a case report [Wilson et al], 4:356
Win, Htet Htet Ne: Charcot-Marie-Tooth disease complicating type 2 diabetes [Win et al], 4:349
Woodburn, James: Reliability and validity of the INFOOT three-dimensional foot digitizer for patients with rheumatoid arthritis [De Mits et al], 3:198
Wrobel, James S.: The economic value of specialized lower-extremity medical care by podiatric physicians in the treatment of diabetic foot ulcers [Carls et al], 2:93
Wukich, Dane K.: The Charcot foot in diabetes [Rogers et al], 5:437

X

Xu, Wen-Jian: Pigmented villonodular synovitis of the ankle: radiologic characteristics [Hao et al], 3:252

Y

Yapar, Nur: Acute phase reactants predict the risk of amputation in diabetic foot infection [Akinci et al], 1:1
Ye, Tianwen: Management of grade III open dislocated ankle fractures: combined internal fixation with bioabsorbable screws/rods and external fixation [Ye et al], 4:307
Yener, Serkan: Acute phase reactants predict the risk of amputation in diabetic foot infection [Akinci et al], 1:1
Yesil, Sena: Acute phase reactants predict the risk of amputation in diabetic foot infection [Akinci et al], 1:1
Yoho, Robert M.: Navicular stress reactions in runners: a review of evaluation and management of a competitive athlete [Yoho and Wells], 5:447
Yu, Jenny: Prevalence of plantar verrucae in patients with human immunodeficiency virus infection during the post–highly active antiretroviral therapy era [Johnston et al], 1:35
Yuan, Wen: Management of grade III open dislocated ankle fractures: combined internal fixation with bioabsorbable screws/rods and external fixation [Ye et al], 4:307

Z

Zhang, Jian-Zhong: Pigmented villonodular synovitis of the ankle: radiologic characteristics [Hao et al], 3:252
Zugaib, Marcelo: Changes in the plantar pressure distribution during gait throughout gestation [Ribeiro et al], 5:415
Zwick, Ernst B.: Lengthening of the midfoot in a case of lower-extremity hemimelia [Kraus et al], 5:456